

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

Z Seeds, TEC and Rutgers, The State Univ. of New Iersen

ALCCRIS, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLED WITH, AND THE TITLE THERETO IS, FROM THE REGORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY THARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLEMISIONEMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE UGHL TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR PRING IT, OR EXPORTING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE PURPOSES, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT BY THE PLANT VARIETY PROTECTION ACT. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

FESCUE, TALL

'Avenger'

In Testimony Whereof, I have hereunto set my hand and caused the seal of the Plant Unriety Protection Office to be affixed at the City of Washington, D.C. this twenty-ninth day of November, in the year two thousand and soven.

Atlast:

Banza

Commissioner
Plant Variety Protection Office
Agricultural Marketing Service

Secretary of

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY - PLANT VARIETY PROTECTION OFFICE	The following statements are made in accordance with the Privacy Act of 1974 (5 U.S.C. 552a) and the Paperwork Reduction Act (PRA) of 1995.		
APPLICATION FOR PLANT VARIETY PROTECTION CERTIFIC (Instructions and information collection burden statement on reverse)	Application is required in order to determine if a plant variety protection certificate is to be issued		
Z Seeds, LLC & Rutgers, The State Univ (31/2006) of New Jersey	v. 2. TEMPORARY DESIGNATION OR 3. VARIETY NAME Roberts LAZ or Avenger		
· ADDRESS (Sireel and No., or R.F.D. No., City, State, and ZIP Code, and Country) • O. Box 8	5. TELEPHONE (Include area code) 410-641-3752 PVPO NUMBER PVPO NUMBER		
TTO 2 Data a rred	reporated, give of incorporation of incorporation syland Jan. 2000		
ohn Zajac, President Seeds, LLC O.Box 8 erlin, MD 21811 William A. Adelphia B 594 Halls Freehold,	Meyer, Ph.D. Research Center Mill Rd. Research Center		
	ajacjj@earthlink.net Tall fescue		
Woodtagoo camaa Birana	15. FAMILY NAME (Botanical) ORCERE 17. IS THE VARIETY A FIRST GENERATION HYBRID? ORCERE NO		
12. CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUBMITTED (Follow instructions of reverse) a. Exhibit A. Origin and Breeding History of the Variety 5. Exhibit B. Statement of Distinctness c. Exhibit C. Objective Description of Variety d	19. DOES THE OWNER SPECIFY THAT SEED OF THIS VARIETY BE SOLD AS A CLASS OF CERTIFIED SEED? See Section 83(a) of the Plant Variety Protection Act) YES (If Yes, answer items 20 KK NO (If 'no', go to item 22; and 21 below) 20. DOES THE OWNER SPECIFY THAT SEED OF THIS YES NO VARIETY BE LIMITED AS TO NUMBER OF CLASSES? IF YES, WHICH CLASSES? FOUNDATION REGISTERED CERTIFIED (BT: 10/31/2006)		
Voucher Sample (2,500 viable untreated seeds or, for tuber propagated varieties, verification that tissue culture will be deposited and maintained in an approved public repository) \$ 3650. Filting and Examination Fee (\$2,705), made payable to "Treasurer of the United States" (Mail to the Plant Variety Protection Office)	21. DOES THE OWNER SPECIFY THAT SEED OF THIS YES NO VARIETY BE LIMITED AS TO NUMBER OF GENERATIONS? IF YES, SPECIFY THE FOUNDATION REGISTERED CERTIFIED NUMBER 1.2.3. etc. (If additional explanation is necessary, please use the space indicated on the reverse.)		
22. FAS THE VARIETY (INCLUDING ANY HARVESTED MATERIAL) OR A HYBRID PRODUCED FROM THIS VARIETY BEEN SOLD, DISPOSED OF, TRANSFERRED, OR USED IN THE U. OTHER COUNTRIES? YES NO FYES. YOU MUST PROVIDE THE DATE OF FIRST SALE, DISPOSITION, TRANSFER, OF FOR EACH COUNTRY AND THE CIRCUMSTANCES. (Please use space indicated on rever	YES X NO		
so, a repail brobadases yasiash a sizena contra mili da performad ni. a begile sabozitosh stile wat	plant variety, and believe(s) that the variety is new, distinct, uniform, and stable as required in Section 47. Act.		
SIGNATURE OF OWNERS	SIGNATURE OF OWNER JAJAN		
Keith R. Cooper	John Zajao		
of Agriculture & Natural SEL TOTAL STREET BY THE SIGNED BY THE Plant Variety Protection Office with WordPariect 60a. Replaces S	capacity or fittle President 5/3/4		

Origin and Breeding History of Avenger (L1Z) Tall Fescue

Avenger (L1Z) tall fescue (*Festuca arundinacea* Schreb.) is a medium low-growing, dark green, medium-fine-leaved, turf-type tall fescue selected from the maternal progenies of 174 clones. Avenger was selected for high shoot density, dark-green color, semi-dwarf growth habit, and medium-early maturity. Approximately 92% of the parental germplasm in Avenger contain the Neotyphodium endophyte.

The 174 parents of Avenger were selected from maternal sources evaluated in progeny turf plots at the Rutgers Plant Science Research and Extension Farm at Adelphia, NJ from the 1997, 1998 and 1999 trials. Thirty-four percent of the maternal germplasm traces to several plants selected from or related to Apache. Approximately fifteen percent traces to a few plants selected from or related to Gazelle. Another fifteen percent traces to a few plants selected from the University of GA State Hospital in 1977. Approximately ten percent traces to a plant selected from the Princeton University campus and used in the development of Rebel. Another six percent traces to plants selected from or related to Amigo. Five percent traces to several plants used in a inter-specific crossing program with perennial ryegrass. Four percent traces to a few plants selected from a farm in North Carolina, 1975 and used in the development of Rebel II. Another four percent traces to a few plants selected from southern GA. Two percent traces to a few plants selected from a park in Lexington, KY in 1979. Another two percent trace to plants selected from Atlanta, GA near GA tech before 1980.

All of the parental germplasm sources of Avenger tall fescue originate from plants selected from old turfs of the United States in a germplasm collection program initiated in 1962, to plants selected from or related to Rebel tall fescue (Funk et al., 1981). Attractive clones were selected from old turfs in Birmingham, AL; Athens, Atlanta, and Millegeville, GA; Preston, ID; Baltimore, MD; Bayonne, Jersey City, Elizabeth, Princeton, and Cape May, NJ; eastern North Carolina; Philadelphia, PA; Nashville, TN; Lexington, KY;

Cincinnati, OH; Dallas, TX; and northern Mississippi. The tall fescue plants selected from old turfs were of unknown origin. All were large patches of turf surviving in stressful environments indicating that they had persisted and developed over a period of many years.

A few hundred attractive, turf-type plants were collected and established in spaced-plant nurseries and/or frequently mowed clonal evaluation trials at Rutgers University. All but a few dozen of the most promising plants were quickly discarded. The best selections were very different from any tall fescue variety in existence at the time of collection. They produced lower-growing turfs with finer leaves, greater density, darker color, and greater tolerance of close mowing.

The most promising plants were identified by their persistence and appearance in old turfs and their performance in spaced-plant nurseries, mowed clonal evaluation tests, and single-plant progeny trails under turf maintenance. Intercrosses of the best performing plants were subjected to varying cycles of phenotypic and genotypic selection depending on their date of collection. New sources of germplasm were added to the breeding program as it became available from the continuing collection program. Each cycle of selection showed continued progress in producing lower-growing, darker green, attractive plants with improved turf performance scores. Selection was also effective in maintaining high seed yields, and good stress tolerance. Substantial progress was made in developing tall fescues with finer leaves, a lower growth profile, increased persistence under close mowing, and increased density.

Large numbers of single-plant progenies were seeded in turf evaluation trials at the Plant Science Research Farm at Adelphia, NJ in 1997, 1998 and 1999. The plants selected for progeny evaluation were selected from spaced-plant nurseries at Adelphia following varying cycles of phenotypic and genotypic selection of germplasm selected from old turfs and germplasm selected from or related to Rebel tall fescue.

Approximately, 7800 tillers were selected from the best performing single-plant progeny turf plots from the 1997, 1998 and 1999 tall fescue test at Adelphia. Eighty-three different single-plot progenies were selected from 1055 plots from 10 different populations from the 1997 test, 635 plots from 9 different populations from the 1998 test and 890 plots from 8 different populations from the 1999 test. These plants were established in greenhouse flats prior to their transfer to a spaced-plant nursery in the spring of 2000 consisting of 3900 plants. Selection was based on performance records as well as appearance at the time the plants were selected from these progeny plots. Selection of plants from each progeny was based on an attractive dark green color, medium-fine leaves, abundant tillering, high shoot density and freedom from disease.

In the spring of 2001, 194 plants were selected from this nursery based on dark green color, semi-dwarf growth habit, high seed yield potential, medium-early maturity and freedom from disease. These plants were moved to an isolated crossing block and inter-pollinated. Twenty-two plants were not harvested due to poor vigor and/or poor floret fertility. Seed from the remaining 174 plants was harvested and designated as breeder seed of Avenger (L1Z) tall fescue. These 174 plants were from 52 different single-plot progenies. These plants produced approximately 27 pounds of breeder seed. Replicated turf plots of Avenger were established at Adelphia in the fall of 2001 and entered in the 2001 National Tall Fescue test to be tested throughout the country. Eleven pounds of breeder seed was sent to Pickseed West for Mr. John Zajac, Berlin, MD. for foundation and certified seed increase.

References

Buckner, Robert C., Jerrell B. Powell, and Rod V. Frakes. 1979. Historical Development, in Buckner, Robert C., and Lowell P. Bush (editors) Tall Fescue. Agronomy Monograph 20. American Society of Agronomy, Crop Science Society

- of America, Soil Science Society of America, Inc., Publishers. Madison, Wisconsin pages 1-8.
- 2. Funk, C.R., R.E. Engel, W.K. Dickson, and R.H. Hurley. 1981. Registration of Rebel tall fescue. Crop Sci. 21:632.

Diagram of Origin and Breeding History of Avenger (LIZ) Tall Fescue

1. 1962 to 1997

Germplasm collection, evaluation, and genetic improvement.

2, 1997-1999

Planted single-plant progenies of plants selected from current cycles of population improvement programs in closely mowed turf trials at Adelphia and North Brunswick, NJ.

3.2000

Selected 7800 plants from 83 of the best performing single-plant progeny turf plots planted in 1997,1998 and 1999. Established 3900 selected plants in an isolated spaced-plant nursery at Adelphia, NJ.

4, 2001

One hundred, ninety-six plants were selected from this nursery and moved to an isolated crossing block. The plants were selected for dark green color, semi-dwarf growth habit, high seed yield potential, medium-early maturity and freedom from disease. Twenty-two plants were not harvested due to poor vigor and/or poor floret fertility. The 174 plants remaining were harvested as Avenger (LIZ) tall fescue. These 370 plants were from 52 different single-plot progenies. These plants produced approximately 27 pounds of breeder seed. Replicated turf plots of Avenger were established at Adelphia in the fall of 2001 and entered in the 2001 National Tall Fescue test to be tested throughout the country. Eleven pounds of breeder seed was sent to Pickseed West/Roberts Seed Co. for foundation and certified seed increase. Additional breeder seed has been produced along with foundation/registered/certified classes of seed for the 5 years between 2001-2006. *Avenger* has been observed to be uniform and stable during this time period. No variants have been observed during this time.

20000002

Exhibit B Summary of Variety Distinctness Avenger Tall Fescue

Avenger tall fescue is a new cultivar. Avenger is most similar to the cultivar Silverado. The two cultivars can be distinguished based on anthesis date. Avenger flowers 6 days earlier than Silverado (Table 1).

Table 1. Anthesis date of tall fescue cultivars for two growing seasons in western Oregon.†

Cultivar	2002	2003	·
Avenger	June 2	May 28	
Watchdog	May 31	May 26	
Kentucky 31	June 2	May 26	
Bonsai	June 6	June 2	
Wrangler	June 4	May 29	
Crewcut	June 7	June 1	
Bonanza	June 6	June 1	
Silverado	June 8	June 3	
LSD@0.05	1 day	1 day	

[†] Data collected from spaced planted progenies of each cultivar from a nursery established at the research facility of Pickseed West, Inc., Albany, OR. Nursery was established in November 2001, using randomized complete block experimental design, with three replications of 20 progeny for each replication. Progeny were observed both for the 2002 and 2003 seasons.

U.S. DEPARTMENT OF AGRICULTURE PLANT VARIETY PROTECTION OFFICE, AMS, USDA NATIONAL AGRICULTURAL LIBRARY Bldg., Rm. 500 10301 BALTIMORE Blvd. **BELTSVILLE, MD 20705**

OBJECTIVE DESCRIPTION OF VARIETY TALL & MEADOW FESCUES

(Festuca spp.)

(2 0 0	once off.)	
NAME OF APPLICANT(S) Z Seeds, and Rutgers, The State of New Jersey	TEMPORARY DESIGNATIO	N VARIETY NAME Avenger
BT:10/3/2006)	l	1
ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP Code) P.O. Box 8, Berlin, MD 21811 Cook College, 88 Lipman Dr., No.	ew Brunswick, NJ 08901-8525	FOR OFFICIAL USE ONLY PVPO NUMBER
		200400273
Place the appropriate number that describes the varietal characteristic of Characteristics described, including numerical measurements, should a SPACED PLANTS. Royal Horticultural Society or any recognized cold asterisk * are characteristics which should be recorded.	represent those that are typical for or fan may be used to determine pl	the variety. Measured data should be fo
* 1. SPECIES: (With comparison varieties, use varieties within the spec	cies of the application variety)	
11 = F. arundinacea (Tall)	<u>Cypes</u>	
1 = Kentucky 31 2 = Rebel 3 = Olympic 7 = Shortstop 8 = Silverado 9 = Rebel Jr. 13 = Caynne		Arid 6 = Rebel II = Crewcut 12 = Bonsai
<u>Forag</u>	e Types	
20 = Kentucky 31 21 = Martin 24 = Kenhy 25 = AU Trium	22 = Forager 23 = Mozark ph26 = Fawn 27 = Cajun	
2 = F. pratensis (Meadow)		
30 = Admira $31 = Beaumont$ $32 = C$	Comtessa 33 = Ensign 34 =	= Trader
* 2. CYTOLOGY:		
Chromosome Number		
3. ADAPTATION: (0 = Not Tested; 1 = Not Adapted; 2 = Adapted)		
2Transition Zone2West2Norther	ast2 Other (Specify):Paci	fic region
* 4. MATURITY: (Date First Headed, 10% of Panicle Emergence)		
4 Maturity Class	h 3 = Early (Fawn) 4 = ado) 8 = 9 =	K31, Kenhy 5 = Medium (Rebel) Very late
Date HeadedMay 5 Location	onAlbany,OR	
5 Days earlier than 4		
——————————————————————————————————————		4
Maturity same as1_ Comparison Variety		9.

	73.1 cm Heig	ht	11.9 cm Internode length				
	13.7 cm shorter t	than4_	3.2 cm shorter than4				
	Height same as	_8 Comparison Va	ariety Length same as _8_ Comparison variety				
	cm taller than	MANAGEMANA	cm longer than				
Æ	IGHT AT EAR EMERGEN	NCE CM: (Flag leaf height	from crown to flag leaf node)				
	32.3 cm Height						
		an4_					
	Height same as	_8 Comparison Va	riety				
	cm taller than						
GI	ROWTH HABIT: (Mature P	Plants)					
) 3 = Semiprostra Rebel) 9 = En	te () 5 = Horizontal () ect (Mini Mustang)				
RI	IIZOMES (Psuedo):						
	mm Length	$X_1 = Absent()$	2 = Rare (Rebel) $3 = Common ($				
LE	mm Length AF BLADE: (Tiller leaves/		2 = Rare (Rebel) 3 = Common ()				
LE	AF BLADE: (Tiller leaves/ *_5 Color: 1 = Lig 7 = Me	turf color)	3 = Medium light green () 5 = Green () 9 = Very dark green ()				
LE	*_5 Color: 1 = Lig 7 = Me5 Specify	turf color) cht green () cdium dark green ()	3 = Medium light green () 5 = Green () 9 = Very dark green () ty				
LE	#_5 Color: 1 = Lig 7 = Me5 Specify *_1_ Anthocyanin:	turf color) tht green () dium dark green () rating of comparison varie	3 = Medium light green () 5 = Green () 9 = Very dark green () ty 9 = Present ()				
LE	*_5 Color: 1 = Lig 7 = Me5 Specify *1_ Anthocyanin: *_9 Basal Hairs:	turf color) tht green () dium dark green () rating of comparison varie 1 = Absent ()	3 = Medium light green () 5 = Green () 9 = Very dark green () ty 9 = Present ()				
LE	*_5 Color: 1 = Lig 7 = Me5 Specify *_1_ Anthocyanin: *_9 Basal Hairs: *_5 Margins:	turf color) tht green () dium dark green () rating of comparison varie 1 = Absent () 1 = Absent ()	3 = Medium light green () 5 = Green () 9 = Very dark green () ty 9 = Present () 9 = Present () 5 = Semi-rough () 9 = Rough ()				
	*_5 Color: 1 = Lig 7 = Me5 Specify *_1_ Anthocyanin: *_9 Basal Hairs: *_5 Margins:	turf color) tht green () dium dark green () rating of comparison varie 1 = Absent () 1 = Smooth () 1 = Very coarse () 7 = Fine ()	3 = Medium light green () 5 = Green () 9 = Very dark green () ty 9 = Present () 9 = Present () 5 = Semi-rough () 9 = Rough () 3 = Coarse () 5 = Medium () 9 = Very Fine ()				
	*_5 Color: 1 = Lig 7 = Me5 Specify *1_ Anthocyanin: *_9 Basal Hairs: *_5 Margins: *_5 Width Class:	turf color) that green () dium dark green () rating of comparison varie 1 = Absent () 1 = Absent () 1 = Smooth () 1 = Very coarse () 7 = Fine () First leaf subtending the flag	3 = Medium light green () 5 = Green () 9 = Very dark green () ty 9 = Present () 9 = Present () 5 = Semi-rough () 9 = Rough () 3 = Coarse () 5 = Medium () 9 = Very Fine ()				
	*_5 Color: 1 = Lig 7 = Me 5 Specify *_1_ Anthocyanin: *_9 Basal Hairs: *_5 Margins: *_5 Width Class: ER LEAF LENGTH CM: (F	turf color) the green () dium dark green () rating of comparison varie 1 = Absent () 1 = Absent () 1 = Smooth () 1 = Very coarse () 7 = Fine () First leaf subtending the flag	3 = Medium light green () 5 = Green () 9 = Very dark green () ty 9 = Present () 9 = Present () 5 = Semi-rough () 9 = Rough () 3 = Coarse () 5 = Medium () 9 = Very Fine () teleaf) * TILLER LEAF WIDTH MM:				
	*_5 Color: 1 = Lig 7 = Me5 Specify *_1_ Anthocyanin: *_9 Basal Hairs: *_5 Margins: *_5 Width Class: ER LEAF LENGTH CM: (F13.0 cm Tiller Le	turf color) the green () dium dark green () rating of comparison varie 1 = Absent () 1 = Absent () 1 = Smooth () 1 = Very coarse () 7 = Fine () First leaf subtending the flag	3 = Medium light green () 5 = Green () 9 = Very dark green () y = Present () 5 = Semi-rough () 9 = Rough () 3 = Coarse () 5 = Medium () 9 = Very Fine () teaf) * TILLER LEAF WIDTH MM:				

		*
FLAG LEAF LENGTH CM:	FLAG LEAF WIDTH MM:	
10.4 cm Flag Leaf Length	2.8 mm Flag Leaf Width	
3.3 cm shorter than12	0.6 mm narrower than _1	
Length same as11 Comparison Variety	Width same as12	Comparison variety
cm longer than	mm wider than	· ·
* 9. LEAF SHEATH: (Basal Portion)		
*_9 Anthocyanin (seedling): 1 = Absent (K31)	9 = Present ()	
* 9 Auricle Hairiness: 1 = Absent ()	9 = Present ()	
* 10. PANICLE: (At seed maturity except where noted.)	4	
*_5 Shape: 1 = Narrow-tapering () 5 = Ov	vate () $7 = Oblong ()$	9 = Other (specify)
*_5 Type: 1 = Compact (appressed) 5 = Int	termediate () 7 = Open ()	9 = Other (specify)
*_9 Orientation: 1 = Nodding () 9 = Ere	ect ()	
*_9 Branch Pubescence: 1 = Glabrous ()	9 = Pubescent () 2003 Data	
*_6 Anther Color (At anthesis): 1 = Yellowish Green Main	2 = Green 3 = Bluish Green 5 = Reddish 6= Other (Yellow)	16% yellow-green 10% green
*_3_ Glume Color (At anthesis): 1 = Yellowish Green 4 = Purplish	2 = Green 3 = Bluish Green 5 = Reddish 6= Other (Specify)	74% yellow
*16.9 cm Panicle Length (from base to tip, if nodding,	, straighten; after anthesis)	
3.1 cm shorter than11		
Length same as13 Comparison Variety		
cm longer than		
* 11. SEED: (With Lemma & Pelea)		
*2790 mg per 1000 seeds		
510 mg less than1		
Weight same as11 Comparison Variety		
mg more than		
PALEA: (Keels or Margins) Hairs: 1 = Absent ()	5 = Short (Missouri 96)	9 = Long ()
LEMMA: Hairs: 1 = Absent (Ken	nhy) 5 = Several ()	9 = Many (Missouri 96)
5.7 mm Lemma Length (Mature)	14.6 mm Lemma width (of 10 seeds)	
0.7 mm shorter than11	1.5 mm narrower than _1	
Length same as _8_ Comparison Variety	Width same as13 Comp	arison variety
non temperather	seems and it am the are	

	*AWNS:	_9A	WNS: $1 = A$	besit () 9 = Present (F	alcon)	100	200400273
•	2.3 mm	Awn length	Of those present	L)				
	1.0 mm	Shorter than	_8 (2003	data)				
	Length same	as	_4 Comp	arison V	ariety			
	mm Lo	nger than						
12. DISE	EASE, INSEC	T, AND NE	MATODE REAG	CTION: (0= Not Tested 1=	Least Res	istant 9= M	ost Resistant)
	0_ Melting	-out <i>Drechsl</i>	era poae		_0	Blind See	d <i>Gloeotinia</i>	temulenta
	0_ Leaf Sp	ot <i>D. siccans</i>	•		_0	Dollar Sp	ot <i>Lanzia, M</i>	follerdiscus spp.
	_0 Net Blo	tch <i>D. dictyo</i>	ides		_0	Stem Rust	t Puccinia gr	raminis
	_8 _ Brown Pa	atch <i>Rhizocto</i>	onia solani		_0	T. Blight i	Typhula inca	urnata
	_0 C. Leaf	Spot <i>Cercos</i>			_0	Pythium E	Blight <i>Pythiu</i>	m spp.
	0_ Pink Sno	ow Mold <i>Ge</i>	rlachia nivalis		_0	Powdery N	Mildew <i>Erys</i>	iphe graminis
	_0 Silver T	op <i>F. tricinci</i>	tum, F. roseum		_0	Crown Ru	st <i>Puccinia</i> d	coronata
	_ 8 Other D	iseasePi	nk Patch				· · · · · · · · · · · · · · · · · · ·	-
	Other Inse	ect						
· Nation	Other Ner	matode						
13. ENV	VIRONMENT	AL STRESS						
	_7 Drought	Stress	1 = Susceptible	()	5 = Tolerant () 9 = Re:	sistant ()	
	Shade Str	ess	1 = Susceptible	()	5 = Tolerant () 9 = Re	sistant ()	
<u>.</u>	_8 Winter S	tress	1 = Susceptible	()	5 = Tolerant () 9 = Res	sistant ()	
14. GIVE indicate the	VARIETY One degree of re	R VARIETI	ES THAT MOS	r CLOSE	ELY RESEMBLE	THE APP	LICATION	VARIETY. For the following characteristics
					me as 3 = More t	han hattar	arastar dar	ilrae ata
Character		Varieties		., 2 54	Chara		Varieties F	
					Chara		v at rettes r	Caung
Leaf Widt	h	Bonsai	2		Leaf (Color	Bonsai	2
Panicle Co	olor	Rebel II	2		Panic	le Shape	Bonsai	2
Seed Size		Bonsai	2		Cold 1	Injury	Cayenne	2
Winter Co	olor	Bonsai	3		Heat		Cayenne	2
Disease (F	Brown Patch)		Cavan	na 2				

*15. EXPERIMENTAL: Give a brief summary of the experimental design utilized to collect the data use on the number of plants measured and plant spacing must be expecified.

on this form. Cultural conditions,

200400273

Unless noted otherwise, data supplied for Exhibit C were generated in the 2002 season from a spaced planted nursery of individuals from 32 cultivars (or experimental lines) cultured at the research facility of Pickseed West, Inc., Albany, OR. The nursery was established in November 2001. Treatments were arranged in a randomized complete block experimental design with three replications. Each replication for an entry was represented by 20 individuals, transplanted in a single row 50 cm apart within the row. Plant nutrition followed 39.2 kgN/ha⁻¹ at transplanting and again in October 2002. Additionally, 100.8kgN/ha⁻¹ was split applied in the spring of 2002 and again in 2003. One half of the spring N was applied in March; the other half was applied in April each of the two years.

Scores for items in part 12 of Exhibit C were taken from the 2002 NTEP data progress report no. 03-1.

	Include form number and edition date on al	reproductions. F	ORM APPROVED - OMB No. 0581-0055			
AGRICULT	RTMENT OF AGRICULTURE URAL MARKETING SERVICE EXHIBIT E F THE BASIS OF OWNERSHIP	Application is required in order to determine if a plant variety protection certificate is to be issued (7 U.S.C. 2421). The information is held confidential until the certificate is issued (7 U.S.C. 2426).				
1. NAME OF APPLICAN		2. TEMPORARY DESIGNATION	3. VARIETY NAME			
Z Seeds, LLC and Rut	gers, The State Univ. of New Jersey	OR EXPERIMENTAL NUMBER Roberts L1Z or L1Z	Avenger			
4. ADDRESS (Street and N	o., or R.F.D. No., City, State, and ZiP, and Country)	5. TELEPHONE (Include area code)	6. FAX (Include area code)			
P.O. Box 8 Berlin, MD 21811	Cook College 88 Linman Dr.	(410) 641-3752	(410) 641-3754			
Dami, 1115 21011	New Brunswick, NJ 08901-8525	7. PVPO NUMBER 200400273				
8. Does the applicant ow	n all rights to the variety? Mark an "X" in the	· ·				
9. Is the applicant (indivi	dual or company) a U.S. national or a U.S. b	pased company? If no, give name of co	ountry. YES NO			
10. Is the applicant the o	riginal owner? YES	NO If no, please answer one	of the following:			
a. If the original right	s to variety were owned by individual(s), is ((are) the original owner(s) a U.S. National NO If no, give name of count				
b. If the original righ	ts to variety were owned by a company(ies)	, is (are) the original owner(s) a U.S. bar NO If no, give name of countr				
11. Addițional explanatio	n on ownership (Trace ownership from origii	nal breeder to current owner. Use the re	everse for extra space if needed):			
PLEASE NOTE:						
Plant variety protection ca	an only be afforded to the owners (not licens	sees) who meet the following criteria:				
	ty are owned by the original breeder, that phich affords similar protection to nationals of					
If the rights to the varie nationals of a UPOV m genus and species.	ty are owned by the company which employ ember country, or owned by nationals of a c	yed the original breeder(s), the company country which affords similar protection t	must be U.S. based, owned by o nationals of the U.S. for the same			
3. If the applicant is an ov	vner who is not the original owner, both the	original owner and the applicant must m	eet one of the above criteria.			
The original breeder/own Act for definitions.	er may be the individual or company who di	rected the final breeding. See Section 4	1(a)(2) of the Plant Variety Protection			
control number. The valid OMB of	ction Act of 1995, an agency may not conduct or sponsor, ontrol number for this information collection is 0581-0055, e instructions, searching existing data sources, gathering a	The time required to complete this information collec	tion is estimated to average 0.1 hour per response,			
marital of family status, political be	e (USDA) prohibits discrimination in all its programs and a eliefs, parental status, or protected genetic information. (N ation (Braille, large print, audiotape, etc.) should contact U	lot all prohibited bases apply to all programs.) Persor	is with disabilities who require alternative means for			
To file a complaint of discrimination 720-5964 (voice and TDD). USDA	n, write USDA, Director, Office of Civil Rights, Room 326- is an equal opportunity provide and employer.	W, Whitten Building, 14th and Independence Avenue,	SW, Washington, D.C. 20250-9410 or call (202)			